

Received 28.10.2011
Reviewed 22.12.2011
Accepted 23.01.2012A – study design
B – data collection
C – statistical analysis
D – data interpretation
E – manuscript preparation
F – literature search

Effect of night and day temperature on the cover colour and quality parameters of apple

Laszlo LAKATOS^{1) ABCD}, Małgorzata BINIAK-PIERÓG^{2) EF},
Andrzej ŻYROMSKI^{2) EF}

¹⁾ University of Debrecen, Centre for Agricultural Sciences and Engineering, Department of Agrotechnology, Böszörményi str. 138, 4032 Debrecen, Hungary, tel. +36-52-508-325 e-mail: lakatos@agr.unideb.hu

²⁾ University of Environmental and Life Sciences, Institute of Environmental Development and Protection, pl. Grunwaldzki 24, 50-363 Wrocław, Poland, tel. +48 71 320-55-69, e-mail: andrzej.zyromski@up.wroc.pl

Abstract

The colour of fruits is considered to be an important quality indicator. Sale ability greatly depends on how well covered the colour is of the specific type of fruits. We used a visual colour scale (0–100 percent) for evaluation of colorization. We distinguished the yellow, green and red cultivars. In the following studies we show those results which consider the red type apples only.

Cover colour is one of the phenometric variables and it is a well-known fact that significant differences can be experienced year by year. The experienced oscillation can be the cause of inappropriate water- and nutrient supply; however it can be the result of some kind of plant disease, extremely high or low temperature, setting rate above the average and outstanding fruit density.

In the present examination it is postulated that the degree of cover colour is mostly influenced by day and night temperature. Therefore, our study aims to find out whether it is true or not. Cover colour belongs to those phenometric characteristic features, only the final value of which is taken into consideration; due to their nature of establishment or forming time it seems useless to follow closely the change in the time of vegetation.

We analysed the ripening groups of apple varieties separately. We have found that at winter ripening cultivars has bigger importance of genetic properties whilst at summer and autumnal ripening ones the weather effects have bigger importance.

The research results shows that the day and night temperature has an important role on fruit quality parameters as sugar content, vitamin C content and on acidity also. Because we can modify the canopy temperature by using evaporative cooling system we have a chance to set up an optimal value of day and night temperature difference in the orchards.

Key words: *cover colour of apple fruits, fruit quality, night and day temperature*

LITERARY OVERVIEW

In the '50s ZERINVÁRI [1950], BERÉNYI and JUSTYÁK [1956] started the survey in orchards as well as in mountainary vineyards. The first results of meteorological examinations on vineyards and orchards were published in the '60s due to the researches of BOGNÁR and KOZMA [1961]. The analy-

ses of macro and microclimatic impacts on the growth of fruits is linked with the name of SZÁSZ [1961]. At the end of the sixties studies analysing the mutual effect of phenological, stage and meteorological parameters were published, CSÖBÖNYEI and STOLLÁR [1969] mainly studied apple and grape cultures. These studies meant great help to develop orchard crop safety and in finding out about the productivity of

different species. As a result of some really hard winters in the decade, the first studies on frost protection [PLETSEK, RADNAI 1964] seemed to come out. In the '70s, several research results were published revealing the relations between fruit growing, dry matter content and weather conditions. Similarly, during this period evapotranspirational researches came to light [FÜRI, KOZMA 1975], as well as studies about apple's water consumption [GERGELY, STOLLÁR 1978], and results relating to Jonathan apple's ripening period estimated by weather variables [STOLLÁR 1977]. In the '80s, greater and greater emphases were laid choosing the production site by means of the description of meteorological background of different varieties [STOLLÁR 1984; STOLLÁR, ZÁRBOK 1981], temperature and radiation state of stands [DUNKEL *et al.* 1981] and examinations of critical winter temperature on surviving winter, analysing mainly grapes [CSAPÓ 1984; DUNKEL, KOZMA 1981].

MATERIAL AND METHOD

The examination material used derives from the assortment of Újfehértói Gyümölcsstermesztési Kutató és Szaktanácsadó Kht (Fruit growing Research and Consultant non-profit company, Újfehértó).

In the examination we recorded and measured 2 trees/varieties in repetition system, the phenological phases and phenometric indicator of 586 kinds of apples, during the years of 1984–2001. Ripening period according to groups:

- 1) ripening in summer,
- 2) ripening in autumn,
- 3) ripening in winter.

In the sample we separated the varieties with good cover colour and later we analysed reciprocal effect with meteorological factors only in these varieties. We made an examination on what reciprocal effect of average day and night temperature on the degree of cover colour has in different months of the vegetation period and 30 days before ripening. Among the varieties, there were the so-called "old" varieties, that were getting to be squeezed out of cultivation, popular and widespread varieties as well as perspective, spreading varieties. A total number of 1172 trees were examined. The different kinds of apples were planted by MM106 stock grafts on spindle shaped parent branched plantation in 1981–1982. The difference between lines and stocks is 8×2 m.

Observation and surveying took place in the research area of Újfehértó. The position of the territory is plain, 115 m a.s.l., 19 km south of Nyíregyháza. Its soil evolved on sand soil-creating stone non-carbonate several layer humous sand, which has a strong, sour acidity (pH 5.74–5.79). Its organic matter content is low (<1%) in its genetic category.

In the examining period, from microclimatic date, air temperature was set hourly and on daily basis

by means of a computer demodulation automatic meteorological measuring station.

In our calculations, we used the below-mentioned temperature variables:

- night temperature T_{night} ,
- day temperature T_{day} ,
- difference between day and night temperature T_{diff} .

During the records, cover colour degree was set in percentage (0–100%) in intervals.

The evaluation of data was fulfilled by means of Excel 97 for Windows programmes. We calculated average and standard deviation from the details. The correlation between phenometric indicators and meteorological elements were evaluated by linear regression and correlation analysis.

RESULTS, CONSEQUENCES

On the basis of frequency distribution of cover colour values it can be stated that almost half (49.7%) of the varieties ripening in summer has a good cover colour, while 25.2% of this group can be counted to be a less well-colouring variety. Those kinds of apples represent the biggest substrate (32.1%-ot) in the sample, whose colouring rate reaches 64–81%. In the sample the varieties with good cover colour appear in a significantly larger proportion than varieties with weak cover colour.

The kinds of apples ripening in autumn show a fairly stable distribution in respect to colouring. 42.5% of these apples can be characterized with good cover colour, while 36.4% have weak cover colour.

In the sample, the number of varieties with good cover colour and weak cover colour seems to be almost equal.

The frequency distribution of apples ripening in winter can be described by a U-shape function. This means that in the sample we can experience the predominance of well as well as weak colouring varieties. We can notice the smallest proportion of varieties with mid-range cover colour in this group.

If cover colour values are examined in the complete range of varieties, as a whole, the following can be stated: 34.1% of the examined varieties can be described with a colouring rate above 70%. Hereinafter, these apples will be considered well colouring varieties (Fig. 1). In case of 24.6% of the sample, colouring rate was between 10–39%. These apples belong to the weak-colouring category. The rate of cover colour in case of 41.2% of the assortment turned out to be between 40–70%; they can be regarded as mid-range colouring varieties.

As a next step, in case of well colouring varieties, we examined how day and night temperature and the difference of these temperatures influence the degree of the fruit's cover colour.




Fig. 1. Distribution of apple cover colour at 586 apple cultivars

According to our hypothesis, the degree of cover colour is mostly influenced by day and night temperature.

In the last 20 years the average night temperature of growing season has increased greatly in the examined habitat. Higher night temperature generally has an unfavourable result on production. Respiration becomes more intense and; therefore, daily weight growth becomes less.

In case of higher night temperature several quality indicators, e.g. sugar content also reaches a lower value, since the energy necessary because of the increased respiration is covered by the plant's own reserves.

Naturally it is also important to examine what changes characterize day temperature in time during the same period. If day temperature increased more intensely, there is no need to worry, because all climatic conditions are given for growth. Even the same rate of growth in day and night temperature does not cause any problem. Problems can occur from the growers point of view if the two examined variables' rate of growth is different.

According to the results, the day temperature's rate of growth falls behind the same value of night temperature.

If we take the difference of day and night temperature and analyse what change the time series of this value shows in the examined period, the following statements can be done:

In the last 20 years the difference of average day and night temperature characteristic to the growing season has decreased, which can be considered significant in 5% rate (Fig. 2). The decreasing trend proves that the increase rate of night temperature exceeds the change of day temperature. This means that if this tendency continues the same way, further decline in crop quality can be expected.

In the next parts of the study we show how the cover colour of apples is influenced by night and day temperatures characteristic to different months and their differences. Night temperature in August in $p = 5\%$ level show a significant relation with the cover colour of apples ripening in summer (Fig. 3) According to the results 5 degrees higher night temperature results in a 15–17% lesser degree of cover colour.




Fig. 2. Time series of the average night and day temperature difference (Újfehértó, between 1984–2001)




Fig. 3. Relationship between night temperature and apple cover colour in August at summer ripening apple varieties

While high night temperature has an unfavourable effect on the establishment of cover colour, in case of higher day temperature a better cover colour can be expected. In case of apples ripening in autumn the relation between day temperature in October and the degree of cover colour is illustrated in Figure 4. The results show that in case of apples ripening in autumn, a 4°C higher temperature results in 10–12% more favourable colouring rate.




Fig. 4. Relationship between day temperature and apple cover colour in October at autumnal ripening apple varieties

In further parts of the study we demonstrate by different ripening groups which month's difference in

day and night temperature has the greatest influence on the cover colour and what function can be used to show the interrelation.

VARIETIES RIPENING IN SUMMER

In case of apples ripening in summer the difference between average day and night temperature in August showed a significant relationship with cover colour degree in 1% rate (Fig. 5).




Fig. 5. Relationship between day and night temperature difference and apple cover colour in August at summer ripening apple varieties

However, the character of the relation cannot be considered linear. That is, in case of big differences between day and night temperatures, a small extent in change does not result in significant change of cover colour. Furthermore, it can be stated that in case of a greater than 6,2°C difference between day and night temperature the extent of cover colour of apples ripening in summer does not increase any more, but it starts decreasing (Fig. 5).

VARIETIES RIPENING IN AUTUMN

In case of varieties ripening in autumn the difference between day and night temperatures in September showed a significant relationship with the cover colour degree in 1% rate (Fig. 6). On the basis of the presented regression relationship it can be stated that if the difference of temperature is increased by




Fig. 6. Relationship between day and night temperature difference and apple cover colour in September at autumnal ripening apple varieties

4°C, as a result, we can reach a 10% higher cover colour degree. By colouring irrigation, this effect can be reached in the years when there is a small difference between day and night temperatures.

VARIETIES RIPENING IN WINTER

In case of apples ripening in winter the difference between average day and night temperature in October showed a significant relationship with cover colour degree in 1% (Fig. 7).




Fig. 7. Relationship between day and night temperature difference and apple cover colour in October at winter ripening apple varieties

The cover colour change as a result of temperature difference could be observed in the smallest extent in case of this group. It can be stated that in case of a greater than 6,2°C difference between day and night temperature the extent of cover colour of apples ripening in summer does not increase any more, but it starts decreasing (Fig. 7).

If the exact dates of the examined varieties' ripening and harvest are known, the relationship with the difference between day and night temperatures might be even closer.

In Figure 8 it is clearly seen that if the difference between the average day and night temperatures increases from 3 to 8°C, in the meantime cover colour degree also increases from 54 to 78%.




Fig. 8. Relationship between apple cover colour and average night and day temperature difference 1 month before ripening at summer ripening apple cultivars in apple gene bank plantation /Újfehértó, 1984–2001


CONCLUSIONS

It is very important to know about day and night temperatures to estimate the extent of fruit cover colour. The presented regression relationships clearly proved that knowing the examined meteorological information the rate of colouring can be estimated

In case of varieties ripening in winter genetical characteristics of the variety have a greater effect, while in case of apples ripening in summer or autumn the year plays a bigger role. „Spúr” varieties usually have a more intense cover colour, since because of the smaller tree size, they are influenced more by weather conditions. In the future it would be useful to analyse how cover colour changes in case of green and yellow varieties, since their market value and saleability are greatly influenced by these factors.

In the future it would also be practical to analyse the character of relationship and interrelation in other habitats with different soil and climatic characteristics. Since factor values received by regression procedures cannot be generalized to other areas without more thorough evaluation.

Acknowledge



REFERENCES

- BERÉNYI D., JUSTYÁK J. 1956. Fenológiai felvételezés hegyvidéki szőlőállományban [Phenological measurements in a mountain vineyard]. *Időjárás*. No 2 p. 104–111.
- BOGNÁR K., KOZMA F. 1961. Együttes szőlőgyümölcsstermesztés mikrometeorológiai vizsgálatáról [Micrometeorological researches in a vineyards and orchards]. *Időjárás*. No 6 p. 366–369.
- CSAPÓ P. 1984. Szőlőültetvények téli fagykárának becslése [Prediction of winter frost damages in vineyards]. *Légtör.* No 1 p. 19–21.
- CSÖBÖNYEI I., STOLLÁR A. 1969. Az alma rügyfakadása és a rügyfakadás-virágzás fenofázis összefüggése a léghőmérséklettel [Relationship between the beginning of blooming length of blooming and air temperature at apple orchards]. *Kísérletügyi Közlemények*. *Kertészet*. No 1-3 p. 19–23.
- CSÖBÖNYEI I., STOLLÁR A. 1969. Kapcsolat a jonathán-alma terméseredménye és az időjárási elemek között [Relationship between the yield of apple (Jonagold) and weather parameters]. *OMSZ. Beszámolók az 1969-ban végzett tud. kut.-ról*. p. 157–161.
- DÁVID A., GERGELY I., STOLLÁR A. 1975. A meteorológia elemek hatása a gyümölcs növekedésére és szárazanyag tartalmára [Effects of meteorological elements on fruit development and dry matter content of fruit]. *OMSZ. Beszámolók az 1975-ben végzett tud. kut.-ról*. p. 150–157.
- DUNKEL Z., KOZMA F. 1981. A szőlő téli kritikus hőmérsékleti értékeinek területi eloszlása és gyakorisága Magyarországon [Spatial distribution of critical low temperature in winter time at grape in Hungary]. *Légtör.* No 2 p. 13–15.
- DUNKEL Z., KOZMA F., MAJOR Gy. 1981. Szőlőültetvényeink hőmérséklet- és sugárzás-ellátottsága a vegetációs időszakban [Temperature and solar radiation supply for vineyards during the vegetation period]. *Időjárás*. No 4 p. 226–234.
- FÜRI J., KOZMA F. 1975. A szőlő tényleges evapotranszpirációja és öntözővíz szükséglete [The actual evapotranspiration and irrigation demand of grape]. *OMSZ. Beszámolók az 1975-ben végzett tud. kut.-ról*. p. 138–145.
- GERGELY I., STOLLÁR A. 1978. Almaültetvények és tenyészvényben nevelt fák vízfogyasztásának vizsgálata [Researching water usage of apple trees]. *OMSZ. Beszámolók az 1978-ban végzett tud. kut.-ról*. p. 138–145.
- JUSTYÁK J. 1985. Szőlőfajták növekedésanalízise Tokajhegyalján [Growth analysis of grape cultivars at Tokaj region]. A klímapotenciál és az agrometeorológiai információk népgazdasági hasznosítása. Bp. p. 337–360.
- NYUJTÓ F. 1965. Gyümölcsstermesztés és agrometeorológia az Alföldön [Investigation the correlation of fruit cultivation and agricultural meteorology in the Hungarian Great Plain]. *Kertészet és szőlészet*. No 15 p. 8–9.
- PLETSEYER J., RADNAI K. 1964. Őszibarack fagyvédelme [Frost protection methods at peach]. *OMSZ. Beszámolók az 1964-ben végzett tud. kut.-ról*. II. rész p. 135–146.
- STOLLÁR A. 1977. A gyümölcsstermesztés agrometeorológiai vonatkozásai a Duna-Tisza közén [Agrometeorological consequences of fruit production at Duna-Tisza region]. *Légtör.* No 4 p. 8–10.
- STOLLÁR A. 1977. A meteorológia elemek hatása a jonathán alma érésére [Effect of meteorological parameters on apple (Jonathan) development]. *OMSZ. Beszámolók az 1977-ben végzett tud. kut.-ról*. p. 214–219.
- STOLLÁR A., ZÁRBOK Zs. 1981. A gyümölcsök optimális termőhelyének elemzése hőmérsékleti adottságok alapján [Analysis of optimal cultivation area of fruit cultivars on the basis of temperature circumstances]. *Légtör.* No 3 p. 15–17.
- SZABÓ Z. 1997. A kedvezőtlen meteorológiai hatások mérséklése [Decreasing the unfavourable meteorological effects in orchards]. In: *Integrált gyümölcsstermesztés [Integrated fruit productions]*. Ed. M. Soltész. Budapest. Mezőgazda Kiadó p. 353–359.
- SZÁSZ G., TÖKEI L. (ed.) 1997. Meteorológia mezőgazdáknek, kertészeknek, erdészeknek [Meteorology for Agricultural-Horticultural and Forestry experts]. Budapest. Mezőgazda Kiadó pp. 772.
- SZÁSZ G. 1961. Makro és mikroklímatis hatások a kőszméte bogycsok növekedésére és beltartalmára [Macro and micro climatic effects on the development and quality parameters of gooseberry]. *Időjárás*. No 5 p. 279–288.
- SZÁSZ G. 1988. Agrometeorológia – általános és speciális [Agricultural meteorology – general and specific]. Budapest. Mezőgazdasági Kiadó pp. 462.
- SZÓKE L., KISS E. 1980. Az időjárás hatása néhány szőlőfajta termésének mennyiségére és minőségére [Weather effect on quality and quantity of some grape cultivars]. *Légtör.* No 3 p. 20–22.
- ZERINVÁRI E. 1950. Növényfejlődési megfigyeléseink a gyümölcsfákon [Observations of fruits development in orchards]. *Időjárás*. No 5–6 p. 154–155.

Laszlo LAKATOS, Małgorzata BINIAK-PIERÓG, Andrzej ŻYROMSKI

Wpływ nocnej i dziennej temperatury na wybarwienie i parametry jakościowe jabłek

STRESZCZENIE

Słowa kluczowe: *dzienna i nocna temperatura, jakość jabłek, wybarwienie jabłek*

Dobre wybarwienie owoców jest jednym z najważniejszych wskaźników ich jakości, ponieważ decyduje ono o możliwości zbytu, a zatem opłacalności uprawy. Różni się ono istotnie w poszczególnych latach, do czego przyczynia się z pewnością niewłaściwe zaopatrywanie w wodę i składniki pokarmowe. Przyczyną mogą być także choroby drzew, ekstremalnie niska lub wysoka temperatura powietrza. W niniejszej pracy przeprowadzono analizę wpływu dziennej i nocnej temperatury na stopień wybarwienia czerwonych odmian jabłek. Materiał badawczy pochodził z Újfehértó (Węgry). Analizy przeprowadzono oddzielnie dla trzech grup jabłek: dojrzewających latem, jesienią oraz zimą. Do analiz wykorzystano dzienne i nocne wartości temperatury powietrza oraz ich dobowe różnice. Stopień wybarwienia owoców oceniano w skali procentowej. Przeprowadzone analizy wykazały, że dla odmian dojrzewających w zimie większe znaczenie mają ich właściwości genetyczne, w przypadku pozostałych większą rolę odgrywają warunki pogodowe. Zarówno dzienna, jak i nocna temperatura powietrza ma istotny wpływ na jakość owoców, tj. zawartość cukru, witaminy C czy kwasowość. Poprzez zwiększenie procesu parowania z koron drzew jabłoniowych można obniżyć ich temperaturę. To działanie pozwala na optymalizację różnic między dziennymi i nocnymi wartościami tego elementu.