

Received 28.10.2011
Reviewed 22.12.2011
Accepted 23.01.2012A – study design
B – data collection
C – statistical analysis
D – data interpretation
E – manuscript preparation
F – literature search

Possibility for modification of microclimate in orchards by using evaporative cooling irrigation

Laszlo LAKATOS^{1) ABCDEF}, Andrzej ŻYROMSKI^{2) EF},
Małgorzata BINIAK-PIEROG^{2) EF}

¹⁾ University of Debrecen, Centre for Agricultural Sciences and Engineering, Department of Agrotechnology, Böszörményi str. 138, 4032 Debrecen, Hungary, tel. +36-52-508-325 e-mail: lakatos@agr.unideb.hu

²⁾ University of Environmental and Life Sciences, Institute of Environmental Development and Protection, pl. Grunwaldzki 24, 50-363 Wrocław, Poland, tel. +48 71 320-55-69, e-mail: andrzej.zyromski@up.wroc.pl

Abstract

Micro irrigation is a horticultural practice mainly used to supply water to the orchard. Nevertheless the micro sprayed irrigation has a powerful influence on fruit microclimatic parameters as temperature and air humidity. By the application we can improve the fruit quality parameters as anthocyanine, C-vitamin, sugar content of the fruits. When the air's temperature is high (about 20°C or higher) the evaporative cooling irrigation significantly decreases the plants' surface temperature and air temperature. The cooling effect is stronger when the air is dryer. The beneficial effect of cooling irrigation is the temperature reduction and frost protection. In March 2010, one month earlier than the expected blooming an irrigation system was established to produce anti-frost treatment and regulate the micro-climate of a apricot, peach, plum, apple and pear orchard which belongs to the University of Debrecen (Hungary). The objective of sprinklers was to cool the air by increasing water evaporation and relative humidity.

The results showed that the water sprayed in the orchard by micro-jet influenced decisively the temperature of the plantation. At higher temperatures (around 20°C), the drop of temperature may attain 5–7°C. A low relative humidity of the air may increase the relative effect. When water was applied at 15 minutes intervals for ten times a day, the temperature could be kept low also in the buds. The beginning of bloom could be delayed for 8–14 day at different fruit species. Blooming dynamics was characterized by a logistic curve in the treated as well as in the control plot. In the treated plot, the curve was steeper than in the control one, in spite of equal temperatures measured in the plots. Under Hungarian climatic conditions, the method was successfully used to delay blooming dates. The main result was the diminution of the frost damage in the spring that assured fruit yields.

Key words: *evaporative cooling irrigation, microclimate modification, delay of blooming*

INTRODUCTION

Micro irrigation is a horticultural practice mainly used to supply water to the orchard. Nevertheless the micro sprayed irrigation has a powerful influence on fruit microclimatic parameters as temperature and air humidity. When the air temperature is high (about 20°C or higher) the evaporative cooling irrigation significantly decreases the plant surface temperature.

The cooling effect is stronger in that case when the air is dryer. By using the cooling irrigation frequently the temperature of the canopy (both bud surface and ambient air temperature) can be decreased so the speed of physiological processes will be slower and will result in later blooming of fruit trees. In the case of early blooming varieties the probability of climatic frost is high and can cause serious damages in orchards. The cooling irrigations beneficial effect comes

forward to the temperature reduction and frost protection.

Below the freezing point the coexistence of water and ice provides for the flowers that the surface temperature remains around 0°C, while the temperature of the surrounding air falls to even – 8°C. Frost protective irrigation is a more wide-spread horticultural practice than cooling irrigation but with the use of cooling irrigation the protection against frost could be significantly escalated.

The dynamics of bloom influenced by the seasonal temperature were examined earlier in apple and sour cherry plantations by LAKATOS *et al.* [2006; 2008; 2009]. The effect of over-tree sprinkler irrigation on colorization by cooling 'Sensation Red Bartlett' pear were analyzed by DUSSI *et al.* [1997]. That authors stated that fruits from cooled trees were more red and less yellow than fruit from noncooled trees, resulting in lower hue values by the middle of the harvestable maturity period in both years of study. Accordingly, cooled fruit should be harvested earlier than noncooled fruit to maintain postharvest quality. Differences between cooled and noncooled fruit with respect to hue, surface blush, and rate of firmness loss were more pronounced in a warm season requiring frequent cooling than in a cooler season.

The latest researches on the topic of microclimate modification in the orchards are trend to analyse the effect of the evaporative cooling irrigation on fruit quality parameters [IGLESIAS *et al.* 2002]. By using micro jets above the canopy at least two times a day the cover colour of fruit will get better [IGLESIAS *et al.* 2005]. Furthermore the evaporative cooling irrigation is suitable for improving the anthocyanine content of the fruit as well [IGLESIAS *et al.* 2008].

The aim of this research work was to investigate the effects of evaporative cooling irrigation on orchard micro-climate parameters. We wanted to know how we can modify the daily distribution of temperature inside the tree canopy. We wanted to analyse the effect of evaporative cooling irrigation on the dynamics of blooming and how many days we can delay the start of blooming at the researched Bosc pear variety.

MATERIALS AND METHODS

The effect of evaporative cooling irrigation on the blooming dynamics of pear of three irrigated and non-irrigated pear trees was analysed. The rate of flowering for each tree by observing 100–100 flowers separately, day by day was calculated. We analysed the effect of micro-spraying on the daily distribution of tree surface temperature on windy nights and days, when the average wind speed was higher than 10 m·s⁻¹. We also investigated the effects of evaporative cooling irrigation on those nights when the relative humidity was lower than 75%, the wind speed was lower than 3 m·s⁻¹ and the temperature was higher

than 10°C, and on those days when the daytime maximum temperature was higher than 20°C and the average wind speed was above 10 m·s⁻¹ separately. Below we introduce our research results for this microclimate modification on changes in blooming times.

Experiments were performed on the Experimental Farm of the Debrecen University, Pállag, Hungary. Measurements refer to a pear plantation with a 5×3 m planting design and trained to open centre system. The pear variety was "Bosc Kobak", planted in 1999.

A special station equipped with thermometers performed measurements with high sensitivity platinum sensors mounted on the branches of the trees near to the buds. The sensors were distributed in the crown at five points, in the centre and on the four cardinal directions on the periphery. The sampling time was 10 minutes for each.

The dynamics of bloom were studied according to the system described by NYÉKI [1980; 1989; 1990; 2002].

Experiments started in March 2010. Irrigation system was established for the purpose of producing anti-frost treatments and regulating the micro-climate. The role of sprinklers was primarily the prevention of frost damage, contrarily in the present program the goal was to cool the air by increasing evaporation and relative humidity

The treatment began nearly one month prior to blooming. The position of the micro-sprinklers was planned at three levels: around the trunks, a few cm above to the soil surface, in the crown region and one half meter above the crown.

The program started the sprinklers every 60 minutes for 15 minutes, water was applied at intervals of 15 minutes for ten times a day from 8 am to 18 pm. The cooling effect was uniform in the whole space of the crown. The effects of the various levels were not observed immediately, because the evaporation was not intense during this period, and the cooling activity was extended to the entire crown. The greatest effect was due to contact-evaporation on the wet surfaces. Measurements were made to register the three levels together.

RESULTS AND DISCUSSION

The evaporative cooling irrigation is a useful method for fruit colorization and for blooming delay. But the over-tree micro sprinkle irrigation has a dangerous effect on plant protection, especially in those regions where the climate is not so dry. Funguses, bacteria spread faster and cause different plant diseases if the air humidity around the tree is constantly high. We should analyse economically the cost of irrigation, risk of plant disease, spraying cost and the advantage of higher fruit quality (better fruit colour, higher sugar content). The efficiency of evaporative

cooling irrigation is high in those areas where the temperature and wind speed is high, the air humidity is low. At humid climatic circumstances, where the annual amount of precipitation higher than 600 mm it is not worse to use it.

During one month period, the temperature of the crown space of the sprinkled trees was 1.5–2.0°C lower than in the non-cooled check (Fig. 1). Around noon, the difference was even 3°C. The smallest difference was in the early morning hours: 0.5–1.0°C.

Although at night time there was no actual solar radiation for an intensive evaporation because of the high wind speed the process was working properly on those nights when the temperature was above 10°C and the average wind speed of the nights was higher than 10 m·s⁻¹ (Fig. 2). The surface temperature at the irrigated trees was 3.0–3.5°C lower than non-irrigated ones thanks to the relatively high temperature and wind speed. At low temperatures and doldrums the evaporation is less intense.

It is well known that those days when the wind speed is lower than 3 m·s⁻¹ the evaporation is not so intensive because the saturated air mass can slowly

flow from the wet surface. The high temperature is suitable for intensive evaporation because the speed of water molecule is higher so its bigger kinetic energy helps the molecule to come out from the water surface easily. Those days when the temperature attained at least 20°C at daytime the effect of irrigation on the daily distribution of temperature was intense. On the basis of low wind speed and high temperature it can be stated, that the highest difference between the irrigated and non-irrigated pear plantation can reach 4°C at noon (Fig. 3).

Not only was the high wind speed able to improve the intensity of evaporation but the relatively low air humidity as well. In lower air humidity cases the unsaturated ambient air can absorb many water molecules easily. So the speed of water molecule is higher in those cases when the relative humidity of the air is low. We analysed separately those nights when the air humidity is not so high (less than 75%) in that nights there were suitable circumstances for evaporation. The highest difference in temperature between the irrigated and non-irrigated pear plantation can reach 3.5°C at midnight (Fig. 4).




Fig. 1. Mean daily temperature in the pear plantation with and without irrigation between March 15 and April 15, 2010




Fig. 2. Effect of irrigation those days when the night temperature ($T > 10^{\circ}\text{C}$) and wind speed ($V_{\text{average}} > 10 \text{ m}\cdot\text{s}^{-1}$) was high between March 15 and May 15, 2010




Fig. 3. Irrigation effect on the daily distribution of temperature those days when the wind speed was lower than $3 \text{ m}\cdot\text{s}^{-1}$ and temperature was higher than 20°C – from March 15 to May 15, 2010




Fig. 4. The effect of night time irrigation on the temperature of the pear plantation on the day when the air humidity was below 75% – 2nd of May, 2010

The most conspicuous effect was experienced on those days when the maximum temperature was above 20°C and at the same time there was an intensive solar radiation and the wind speed was higher than $10 \text{ m}\cdot\text{s}^{-1}$, at daytime the irrigated buds were $6\text{--}7^\circ\text{C}$ cooler than the non-irrigated ones. (Fig. 5). If during the research period there had been same weather circumstances as mentioned above several times, the bloom delaying effects of irrigation would have been more efficient.

On warm days, when the maximum temperature reached 20°C , there was an intensive solar radiation and low wind speed-less than $3 \text{ m}\cdot\text{s}^{-1}$ and at the same time the relative humidity of the air was less than 50% at noon, the daily distribution of temperature showed intensive fluctuation (Fig. 6). By usage of micro sprinklers the bud surface showed a decreasing of relatively high temperature ($2\text{--}2.5^\circ\text{C}$) within short intervals.

Irrigation not only lowered the temperature when it was below 10°C but also moderated its fluctuation. On those days, when the sky was cloudy and the temperature was low, we could not find significant differences between irrigated and non-irrigated trees' surface temperature. Irrigation diminished the process of warming up and cooling down, thus the risk of heat-stress and sunburn can be reduced.

The retarding effect on blooming by cooling irrigation was measured on three selected trees and as a control group on three dry trees. Non-irrigated trees started blooming at April 14–15, whereas the irrigated trees by 9–10 days later. In Figure 7, the rising angle of the sigmoid-curve was less steep. The time between start of bloom and full bloom lasted 17 days instead of 14 days in the check. The slowing down of the dynamics lasted during the whole blooming period. The dry check finished bloom at May 2 whereas the irrigated at May 14, but the whole blooming period was prolonged by 3 days.




Fig. 5. The effect of daytime irrigation on the daily distribution of temperature in pear plantation on a windy day when the average wind speed was higher than 10 m s^{-1} – 29th of April 2010




Fig. 6. The effect of daytime irrigation on the daily distribution of temperature on the day when the wind speed was lower than 3 m s^{-1} , and the relative humidity less than 50% at noon – in pear plantation – 23rd of April 2010




Fig. 7. The effect of irrigation on the blooming dynamic of pear. Between April 14 and May 14, 2010


CONCLUSIONS

The micro-sprinkler irrigation is a suitable method for modifying microclimate in orchards

[DUSSE *et al.* 1997]. The evaporative cooling effect of the irrigation mainly depends on weather circumstances. When the temperature is higher than 10°C there is a significant evaporative cooling effect. If the

wind speed is less than $3 \text{ m}\cdot\text{s}^{-1}$ the evaporation is not so intensive the cooling effect is slower and not so efficient than in those cases when the wind speed is higher than $10 \text{ m}\cdot\text{s}^{-1}$. On windy days in spring time the intensity of evaporative cooling irrigation is $1.5\text{--}2.0^\circ\text{C}$ higher than on those days when the wind speed is below $3 \text{ m}\cdot\text{s}^{-1}$. The relative humidity of the air is also important in the evaporation. On those days when the relative humidity was about 50% at noon the evaporative cooling effects' intensity was $1.0\text{--}1.5^\circ\text{C}$ higher than on those days when the humidity was above 75% at early afternoon. The evaporative cooling irrigation was a useful procedure for delaying the start of blooming of Bosc pear variety. The time delaying can reach 9–10 days. By this method, frost damage probability can be decreased by more than 70%, because climatically the frost occurrence at this time of the year is less by 70% at least.

Acknowledge



REFERENCES

- DUSSI C., SUGAR D., RIGHETTI T., AZARENKO A. 1997. Effects of cooling by overtree sprinkler irrigation on fruit color and firmness in 'Sensation' Red Bartlett pear. Hort-Technology. No 7 p. 55–57.
- IGLESIAS I., SALVIA J., TORGUET L., CABÚS C. 2002. Orchard cooling with overtree microspinkler irrigation to improve fruit colour and quality of 'Topred Delicious' apples. Scientia Horticulturae. No 93 p. 39–51.
- IGLESIAS I., SALVIA J., TORGUET L., MONTSERRAT R. 2005. The evaporative cooling effects of overtree microsprinkler irrigation on 'Mondial Gala' apples. Scientia Horticulturae. No 103 p. 267–287.
- IGLESIAS I., ECHEVERRÍA G., SORIA Y. 2008. Differences in fruit colour development, anthocyanin content, fruit quality and consumer acceptability of eight 'Gala' apple strains. Scientia Horticulturae. No 119 p. 32–40.
- LAKATOS L., SZABÓ T., SZABÓ Z., SOLTÉSZ M., NYÉKI J. 2009. Relation of sour cherry blooming dynamics and meteorological variables. International Journal of Horticultural Science. No 15 (4) p. 17–23.
- LAKATOS L., SZABÓ T., RACSKÓ J., SOLTÉSZ M., SZABÓ Z., NAGY J., NYÉKI J. 2006. Effects of weather characteristics on blooming dates in an apple gene bank plantation between 1984 and 2001. International Journal of Horticultural Science. No 12 (2) p. 37–44.
- LAKATOS L., SZABÓ T., SOLTÉSZ M., SUN Z., WANG Y., SZABÓ Z., NYÉKI J. 2008. Időjárási változók hatása a meggy virágzástamának alakulására [Effects of weather parameters on the length of sour cherry blooming time]. Klíma-21 Füzetek. No 53 p. 60–67.
- NYÉKI J. 1980. Gyümölcsfajták virágzásbiológiája és termékenyülése [Floral biology and fertilization of fruit cultivars]. Budapest. Mezőgazdasági Kiadó pp. 334.
- NYÉKI J. 1989. Csonthéjas gyümölcsűek virágzása és termékenyülése [Blooming and fertilization of stone fruit]. Doktori értekezés – kézirat [Thesis – manuscript]. Budapest. MTA p. 88–110.
- NYÉKI J. 1990. A gyümölcsstermő növények virágzása, megporzása és termékenyülése [Blooming pollination and fertilization of fruit species]. In: Gyümölcsstermesztés [Fruit Production]. Ed. F. Gyuró. Budapest. Mezőgazdasági Kiadó p. 61–90.
- NYÉKI J. 2002. Gyümölcsstermő növények virágzása és termékenyülése, ültetvények fajtatársítása [Blooming and fertilization and the association of cultivars in plantation]. Egyetemi jegyzet [Text book of the University], Debrecen. DE ATC pp. 68.

Laszlo LAKATOS, Andrzej ŻYROMSKI, Małgorzata BINIAK-PIERÓG

Możliwość modyfikacji mikroklimatu w sadach poprzez stosowanie nawodnień

STRESZCZENIE

Słowa kluczowe: modyfikacja mikroklimatu, nawadnianie, opóźnianie kwitnienia

Stosowanie mikronawodnień w praktyce ogrodniczej ma duży wpływ na warunki mikroklimatyczne, tj. temperaturę i wilgotność powietrza. Największą korzyścią ze stosowania mikrozaszczyn jest obniżanie temperatury powierzchni roślin oraz ich ochrona przed przymrozkami. W marcu 2010 r., miesiąc przed oczekiwanym kwitnieniem śliw, w sadzie doświadczalnym, należącym do Uniwersytetu w Debreczynie (Węgry), uruchomiono system nawadniający w celu ochrony roślin przed przymrozkami. Na podstawie otrzymanych wyników stwierdzono, że woda rozpylana w sadzie przez mikro zraszacze zdecydowanie modyfikowała temperaturę powietrza w plantacji. W warunkach wyższej temperatury (ok. 20°C) można uzyskać jej obniżenie nawet o $5\text{--}7^\circ\text{C}$, a efekt ten jest większy, gdy wilgotność powietrza jest mała. Zastosowanie dawek wody przez 15 minut 10 razy w ciągu doby pozwoliło na utrzymywanie takiej temperatury także w pąkach roślin. Początek kwitnienia może być opóźniony nawet o 9–10 dni. W warunkach klimatycznych Węgier metoda ta jest z powodzeniem stosowana do opóźniania kwitnienia, co skutkuje zmniejszeniem niekorzystnego oddziaływania przymrozków wiosennych na plon owoców.