

DOI 10.2478/pjvs-2014-0090

Original article

# Methyl paraben as a sex pheromone in canine urine – is the question still open?

M. Dziecioł<sup>1</sup>, J. Politowicz<sup>2</sup>, A. Szumny<sup>2</sup>, W. Nizański<sup>1</sup>

<sup>1</sup> Wrocław University of Environmental and Life Sciences, Department of Reproduction and Clinic of Farm Animals, 50-366 Wrocław, Pl. Grunwaldzki 49, Poland

<sup>2</sup> Wrocław University of Environmental and Life Sciences, Department of Chemistry, 50-375 Wrocław, C.K. Norwida 25, Poland

## Abstract

The literature concerning the issue of canine sex pheromones includes reports presenting completely conflicting opinions about the chemical composition of the canine urine in the context of semiochemical communication. At present, the predominant report cited by many different authors is the article published in Science in 1979 by Goodwin et al., presenting methyl p-hydroxybenzoate (methyl paraben) as the main canine sex pheromone. While it has been proved that pure methyl paraben lacks semiochemical activity as do commercially available products containing this substance (Eau D'Estrus, Synbiotics, USA), in view of the conflicting published reports the aim of this study was to reevaluate using modern techniques the presence of methyl p-hydroxybenzoate in canine urine during different phases of the ovarian cycle. Ten female dogs of different breeds were used. Urine samples from bitches collected during various stages of the ovarian cycle were examined with using the SPME and GC/MS methods. Methyl paraben was not detected in any of the samples. In conclusion, because of the lack of methyl-p-hydroxybenzoate in the samples examined, the present study confirmed negative opinions on the possibility of this substance playing a crucial role in semiochemical communication during reproduction in dogs (*Canis familiaris*).

**Key Words:** sex pheromones, bitch, GC/MS, methyl p-hydroxybenzoate, methyl paraben

## Introduction

In the 1970's and 80's, there was a large interest in the topic of semiochemical communication in animals (Meredith et al. 1980, Novotny et al. 1984, Novotny et al. 1985, Raymer et al. 1986, Singer et al. 1986). While in the beginning rodents were the main group of animals tested for pheromonal communication, soon other species were investigated including wild and

domestic animals (Brownlee et al. 1969, Izard and Vandenberg 1982, Stevens et al. 1982, Raymer et al. 1986, Nishimura 1991, Jezierski 1992, Grzegorzewski 2006, Dziecioł et al. 2012a, Dziecioł et al. 2013). The rapid development of modern methods dedicated to detection and identification of chemical compounds, including gas chromatography-mass spectrometry (GC/MS), fostered studies into the composition of natural bioactive secretions.

In 1979 Goodwin, Gooding and Regnier published in *Science* the results of their study describing sex pheromones in dogs (Goodwin et al. 1979). According to these authors, methyl p-hydroxybenzoate (methyl paraben, MP) is the main substance present in the urine of bitches in heat, and application of MP into the vulvar region of animals that were out of heat (anoestrus) stimulated males to exhibit mating behavior. This observation gave rise to a hypothesis that the main sex pheromone of bitches is just methyl paraben (Goodwin et al. 1979).

An attempt to verify this hypothesis was made by Kruse and Howard in 1983. In their paper published in the *Journal of Chemical Ecology*, they stated that even though they confirmed the presence of methyl paraben in samples of female dog urine, results of behavioral analysis showed that methyl p-hydroxybenzoate cannot be considered as a key sexual attractant for male dogs (Kruse and Howard 1983). It is also worth noting that in the article published by Schultz et al. (1985) the authors did not even include methyl paraben in the list of substances identified in samples of urine collected from bitches in oestrus.

While all the reports mentioned above on the issue of dog sex pheromones collectively presented contradictory conclusions, and therefore created great confusion in the area of canine sex pheromones, the aim of our work was to attempt once again to investigate the composition of canine urine and ascertain if methyl p-hydroxybenzoate can be considered as a dog sex pheromone. Because since the 1980's the accuracy of laboratory procedures like GC/MS has significantly improved, we decided to focus on confirming the presence of candidate substances in canine urine.

## Materials and Methods

### Location, animals and sample collection

The experiment was conducted in the Clinic of the Department of Reproduction and in the Laboratory of the Department of Chemistry belonging to the Wrocław University of Environmental and Life Sciences, Poland. The study was approved by the Local Ethical Committee.

Samples were collected from bitches belonging to the Experimental Breeding Kennel located at the Department of Reproduction and from patients of the local Clinic of Reproduction. All animals included in the experiment were previously clinically examined and no diseases were found in any animals. Moreover, to eliminate the possibility of disease compromising the results, all urine samples were also tested for kidney, bladder and lower urinary tract diseases. Only

samples from healthy females were used for further examination.

Samples of urine were collected from 10 bitches of different breeds (four beagles, four German shepherds, two golden retrievers) with an average age of 4-6 years. The phase of the oestrous cycle was determined according to the protocol described below. From each female, at least three samples of urine were collected during proestrus and three during oestrus. During these phases, the attractiveness of the females to the males was confirmed by performing tests with experienced male dogs belonging to the Experimental Kennel. In addition, urine samples were also collected from all of the females twice during anoestrus and metoestrus. The urine was collected during morning hours directly into a sterile dipper and then immediately transferred into a sterile plastic container and stored at 2-5°C for a maximum of one day.

Detection of the exact stage of the oestrous cycle was achieved by clinical examination and laboratory tests. During clinical examination, the presence and character of the vaginal discharge, as well as the presence and quality of vulvar oedema and tolerance reflex, were evaluated. Laboratory tests consisted of vaginal cytology and analysis of progesterone concentration in peripheral blood (Kustritz 2005, 2006). Progesterone concentration was determined by commercial radioimmunoassays validated for dog blood plasma (Progesterone Coat-a-Count kit, Diagnostic Products Corporation, Los Angeles, CA, USA) (Srikandakumar et al. 1986). Eight ml of blood were taken by venipuncture from the cephalic vein into heparinised tubes. Plasma was separated 10 min after blood collection by centrifugation for 15 min at 2000 x g. Progesterone concentration was determined on the same day using the afore-mentioned RIA method.

### Sample preparation

An individual standard solution was prepared by dissolving 10 mg of methyl paraben in 10 ml of methanol. Then, a series of five working standard solutions ranging in concentration from 0.005 to 5 µg/ml of methyl paraben were prepared. Solid-phase microextraction (SPME) was used to extract methyl paraben from each standard prior to GC-MS analysis. An appropriate volume (1-10 µl) of methyl paraben solution was added to 10 mL of water with 5 mL of dichloromethane and shaken for 2 min. After centrifugation, the organic phase was collected, dried over anhydrous sodium sulphate and condensed in a stream of nitrogen (Canosa et al. 2006). The residue was exposed to SPME fiber (Divinylbenzene/Car-

boxen/Polydimethylsiloxane) for about 20 min at 60°C. After exposure for 20 min, the SPME fiber was retracted into the needle of the holding syringe, solvent drops attached to the metallic needle were removed using a paper tissue and the fiber was exposed to GC-MS for 5 min (Fei et al. 2011). The capability of the equipment used in the experiment to detect methyl paraben was confirmed during a preliminary study, when MP was added to samples of urine obtained from bitches in anoestrus. Because during this part of the study it was also confirmed that the results of detecting methyl paraben in water and in urine were identical, in the experiment water was used as a matrix. Similar approaches were applied also by other researchers (Goodwin et al. 2008, Arnaiz et al. 2014). Risticvic et al. (2010) also used this SPME method for identification of MP in a much more complex matrix which is sawage water.

### Chromatographic conditions

The chemical composition of the volatile compounds absorbed on the fiber was analysed using a gas chromatograph (GC) coupled to a mass spectrometer (MS), using a Saturn 2000 MS Varian Chrompack with a ZB-1 (Phenomenex) column (30 m x 0.25 µm film x 0.25 mm ID). The MS was equipped with an ion-trap analyzer set at 1508 for all analyses with an electron multiplier voltage of 1350 V. Scanning (1 scan s<sup>-1</sup>) was performed in the range of 39-400 m/z using electron impact ionization at 70 eV. The analyses were carried out using helium as the carrier gas at a flow rate of 1.0 mL min<sup>-1</sup> in a split ratio of 1:20 and the following program: 60°C at the beginning and holding for 3 min; 3°C/min up to 120°C; then 15°C/min to 300°C. The injector and detector were held at 200 and 300°C, respectively. Analyses were carried out using helium as the carrier gas at a flow rate of 1.0 mL min<sup>-1</sup>, in splitless mode in SPME and liquid injection mode.

The compound was identified by using three analytical methods: Kovac indices (KI), GC/MS retention times of authentic chemical-standards (S) and mass spectra of compounds and NIST05 spectral library collection (MS). The retention index standards used in this study consisted of a mixture of aliphatic hydrocarbons ranging from C-5 through C-17 dissolved in methanol.

### Results

For the SPME as well as the liquid injection mode of MP detection in urine, the limit of detection

(LOD) and limit of quantification (LOQ) were determined. The initial parameters were established from the signal-to-noise ratios of 3 and 10, respectively. In the case of SPME, LOD was found to be 0.005 µg/ml for methyl paraben, while liquid injection gave a LOD of 0.05 g/mL.

The SPME approach for MP analysis gave satisfactory results. The LOD level was lower than that reported by Lee et al. (2005) of 0.01 µg/ml (defined for a signal-to-noise ratio of 3) using solid-phase extraction, solvent evaporation and derivatization of the analyses with pentafluoropropionic anhydride. Determination was performed using stationary phase C8 with wavelength 254 nm, with separation using a mobile phase of methanol: water (60:40 w/w). The LOD level after optimization of the HPLC method was determined at 0.035 µg/ml (Imamović et al. 2012). In comparison to other methods for the determination of MP in cosmetics, LODs were higher than those obtained using the SPME method. In all our samples tested, in liquid injection mode as well as SPME we did not find any traces of MP.

### Discussion

Semiochemical signalling is one of the oldest means of communication used by organisms of all taxa. Even though studies dedicated to identification of the active biological compounds in animal secretions have been performed for many years in laboratories all over the world, the results and consequently our understanding of this primitive communication method seem to be still not fully satisfactory (Pageat and Gaulnier 2003).

Analyzing the issue of canine sex pheromones, we can clearly see that the publication by Goodwin et al. (1979) is still the most often cited study (Person 1985, Pageat and Gaulnier 2003, Kustritz 2005, Santos et al. 2013, Wani et al. 2013). Described as a primary dog sex pheromone, methyl paraben has become a component of commercially available products (Eau D'Estus, Synbiotics, USA) recommended to both veterinarians and breeders as a useful tool for male canine sexual stimulation (Kustritz 2005, Kutzler 2005). However, analysis of the literature on the activity of methyl paraben in the context of sexual arousal shows a lack of efficiency of this substance (Kruse and Howard 1983, Tonosaki and Tucker 1985, Dzięcioł et al. 2011). In the original article reporting an effective association between PGF<sub>2h</sub> and methyl 4-hydroxybenzoate prior to electroejaculation in dogs, while the authors obtained better results in a group of males exposed to methyl paraben, they also confirmed that there were no signs of sexual arousal observed in

these males (Santos et al. 2013). Also, while the influence of natural pheromones on physiological parameters (e.g., heart rate) in dogs was previously described by Dzięcioł et al. (2012b), Santos et al. (2013) did not find any differences between groups in heart and respiratory rate or rectal temperature. In this case, a probable explanation for the better results obtained in males injected with PGF $2\alpha$  and stimulated by MP is the stimulating influence of PGF $2\alpha$  (Estienne and Harper 2000, Kozink et al. 2002).

In a report by Goodwin et al (1979), except the methyl paraben other substances like diethyl phthalate (DEP) C<sub>12</sub>H<sub>14</sub>O<sub>4</sub>, dibutyl phthalate (DBP) and heptadecane C<sub>17</sub>H<sub>36</sub> were identified as compounds present in canine urine. All these substances could be easily classified as contaminants of the samples: for example dibutyl phthalate (DBP) is a commonly used plasticizer also used as an ectoparasiticide, and diethyl phthalate (DEP) C<sub>12</sub>H<sub>14</sub>O<sub>4</sub> can be transferred from plastics (also plasticizers) and is often used in cosmetics and fragrances. It is worth noting that methyl paraben (CH<sub>3</sub>(C<sub>6</sub>H<sub>4</sub>(OH)COO) (CAS No. 99-76-3) (E218) is also very often present in our environment. It is a stable, non-volatile compound and except for natural sources (fruits) it has been commonly used as an antimicrobial preservative in foods, drugs and cosmetics for over 50 years (Soni et al. 2002). While the influence of MP on the sexual behavior of dogs has been questioned by many authors, its accidental presence in evaluated samples, similar to other chemicals mentioned above, appears to be highly probable.

Taking into account results obtained during our study and emerging doubts after reading reports describing compounds in the urine of bitches, we conclude that methyl p-hydroxybenzoate cannot be considered as the main canine sex pheromone because it is not a normal component of canine urine. This conclusion is in agreement with our previous results presenting the lack of efficiency of commercially available products containing methyl paraben in creating sexual arousal in male dogs (Dzięcioł et al. 2011). Thus, the question of identification of the canine sex pheromones is still open and the need for further detailed studies is indicated.

### Acknowledgements

This research was supported by statutory research and development activity funds assigned to the Faculty of Veterinary Medicine, Wrocław University of Environmental and Life Sciences, Poland. Authors would like to thank dr Barry Bavister for help in the final preparation of the manuscript.

### References

- Arnáiz E, Moreno D, Quesada R (2014) Determination of volatiles in mouse urine by headspace solid phase microextraction and gas chromatography-mass spectrometry. *Anal Lett* 47: 721-729.
- Canosa P, Rodriguez I, Rubi E, Bollain MH, Cela R (2006) Optimisation of a solid-phase microextraction method for the determination of parabens in water samples at the low ng per litre level. *J Chromatogr A* 1124: 3-10.
- Dzięcioł M, Nizański N, Ochota M, Kozdrowski R, Twardoń J (2011) The usefulness of the synthetic bitch sex pheromones for stimulation of reproductive reflexes in dogs (*Canis familiaris*). *Reprod Domest Anim* 46 Suppl 3: 101.
- Dzięcioł M, Nizański W, Ochota M, Kozdrowski R, Stańczyk E (2012a) Observation on possibility to identify by the stud dogs the signs of the fertile period in bitches. *JAVA* 11: 962-967.
- Dzięcioł M, Stańczyk E, Noszczyk-Nowak A, Nizański W, Ochota M, Kozdrowski R (2012b) Influence of bitches sex pheromones on the heart rate and other chosen parameters of blood flow in stud dogs (*Canis familiaris*). *Res Vet Sci* 93: 1241-1247.
- Dzięcioł M, Nizański W, Stańczyk E, Kozdrowski R, Najder-Kozdrowska L, Twardoń J (2013) The influence of antibiotic treatment of bitches in oestrus on their attractiveness to males during mating. *Pol J Vet Sci* 16: 509-516.
- Estienne MJ, Harper AF (2000) PGF $2\alpha$  facilitates the training of sexually active boars for semen collection. *Theriogenology* 54: 1087-1092.
- Fei T, Li H, Ding M, Ito M, Lin JM (2011) Determination of parabens in cosmetic products by solid-phase microextraction of poly(ethylene glycol) diacrylate thin film on fibers and ultra high-speed liquid chromatography with diode array detector. *J Sep Sci* 34: 1599-1606.
- Goodwin M, Gooding KM, Regnier F (1979) Sex pheromone in the dog. *Science* 203: 559-561.
- Goodwin T, Brown PA, Eggert MS, Evola MG, House SJ, Morshedi RG, Weddell ME, Chen CJ, Jackson SR, Aubut Y, Eggert J, Schulte B, Rasmussen L (2008) Use of automated solid phase dynamic extraction (SPDE)/GC-MS and novel macros in the search for african elephant pheromones. In: Hurst J, Beynon R, Roberts SC, Wyatt T (eds) *Chemical Signals in Vertebrates* 11. Springer, New York, pp 25-35.
- Grzegorzewski WJ (2006) The influence of boar pheromones on the contractile reactivity of the isolated superficial veins of the nose and face in ovariectomized prepubertal gilts and in gilts during sexual maturation. *Pol J Vet Sci* 9: 127-133.
- Imamović B, Šober M, Bečić E (2012) HPLC Determination of Some frequently used Parabens in Sunscreens. *Int J Pharm Teach Pract* 3: 219-224.
- Izard MK, Vandenberg JG (1982) Priming pheromones from oestrous cows increase synchronization of oestrus in dairy heifers after PGF-2 $\alpha$  injection. *J Reprod Fertil* 66: 189-196.
- Jeziński T (1992) The effectiveness of estrus detection in cows by a trained dog. *Anim Sci Pap Rep* 10: 57-66.
- Kozink DM, Estienne MJ, Harper AF, Knight JW (2002) The effect of luteal phase on the training of sexually inexperienced boars for semen collection. *Theriogenology* 58: 1039-45.



- Kruse SM, Howard WE (1983) Canid sex attractant studies. *J Chem Ecol* 9: 1503-1510.
- Kustritz MV (2005) Reproductive behavior of small animals. *Theriogenology* 64: 734-774.
- Kustritz MV (2006) Collection of tissue and culture samples from the canine reproductive tract. *Theriogenology* 66: 567-574.
- Kutzler MA (2005) Semen collection in the dog. *Theriogenology* 64: 747-754.
- Lee HB, Peart TE, Svoboda ML (2005) Determination of endocrine-disrupting phenols, acidic pharmaceuticals and personal-care products in sewage by solid-phase extraction and gas chromatography-mass spectrometry. *J Chromatogr A* 1094: 122-129.
- Meredith M, Marques DM, O'connell RO, Stern FL (1980) Vomeronasal pump: significance for male hamster sexual behavior. *Science* 207: 1224-1226.
- Brownlee RG, Silverstein RM, Muller-Schwarze D, Singer AG (1969) Isolation, identification and function of the chief component of the male tarsal scent in black-tailed deer. *Nature* 221: 284-285.
- Nishimura K, Utsumi K, Okano T, Iritani A (1991) Separation of mounting-inducing pheromones of vaginal mucus from estrual heifers. *J Anim Sci* 69: 3343-3347.
- Novotny M, Schwende FJ, Weisler D, Jorgenson JW, Carmack M (1984) Identification of a testosterone-dependent unique volatile constituent of male mouse urine: 7-exo-ethyl-5-methyl-6,8-dioxabicyclo[3.2.1]-3-octene. *Experientia* 40: 217-219.
- Novotny M, Harvey S, Jemiolo B, Alberts J (1985) Synthetic pheromones that promote inter-male aggression in mice. *Proc Natl Acad Sci U S A* 82: 2059-2061.
- Pageat P, Gaultier E (2003) Current research in canine and feline pheromones. *Vet Clin North Am Small Anim Pract* 33: 187-211.
- Person JR (1985) Mounting Evidence of Paraben Sensitivity in Dogs. *Arch Dermatol* 121: 1107.
- Raymer J, Wiesler D, Novotny M, Asa C, Seal US, Mech LD (1986) Chemical scent constituents in urine of wolf (*Canis lupus*) and their dependence on reproductive hormones. *J Chem Ecol* 12: 297-314.
- Risticovic S, Lord H, Górecki T, Arthur CL, Pawliszyn J (2010) Protocol for solid-phase microextraction method development. *Nat Protoc* 5: 122-139.
- Santos IP, Ramos CL, Ramos JL, Oliveira RF, Cunha IC (2013) Efficient association between PGF2 $\alpha$  and methyl 4-hydroxybenzoate sex pheromone prior to electroejaculation in dogs. *Reprod Domest Anim* 48: 160-164.
- Schultz TH, Kruse SM, Flath RA (1985) Some volatile constituents of female dog urine. *J Chem Ecol* 11: 169-175.
- Singer AG, Macrides F, Clancy AN, Agosta WC (1986) Purification and analysis of a proteinaceous aphrodisiac pheromone from hamster vaginal discharge. *J Biol Chem* 261: 13323-13326.
- Singer AG (1991) A chemistry of mammalian pheromones. *J Steroid Biochem Mol Biol* 39: 627-632.
- Soni MG, Taylor SL, Greenberg NA, Burdock GA (2002) Evaluation of the health aspects of methyl paraben: a review of the published literature. *Food Chem Toxicol* 40: 1335-1373.
- Srikandakumar A, Ingraham RH, Ellsworth M, Archbald LF, Liao A, Godke RA (1986) Comparison of a solid-phase, no-extraction radioimmunoassay for progesterone with an extraction assay for monitoring luteal function in the mare, bitch and cow. *Theriogenology* 26: 779-793.
- Stevens K, Perry GC, Long SE (1982) Effect of ewe urine and vaginal secretions on ram investigative behavior. *J Chem Ecol* 8: 23-29.
- Wani AA, Dhindsa SS, Shafi TA, Chowdhary SR, Kumar B (2013) The role of pheromones in animal reproduction – a review. *Prog Res* 8: 14-18.