

Symmoriiform sharks from the Pennsylvanian of Nebraska

MICHAŁ GINTER

University of Warsaw, Faculty of Geology, Żwirki i Wigury 93, PL-02-089 Warsaw, Poland.

E-mail: m.ginter@uw.edu.pl

ABSTRACT:

Ginter, M. 2018. Symmoriiform sharks from the Pennsylvanian of Nebraska. *Acta Geologica Polonica*, **68** (3), 391–401. Warszawa.

The Indian Cave Sandstone (Upper Pennsylvanian, Gzhelian) from the area of Peru, Nebraska, USA, has yielded numerous isolated chondrichthyan remains and among them teeth and dermal denticles of the Symmoriiformes Zangerl, 1981. Two tooth-based taxa were identified: a falcatid *Denaëa saltsmani* Ginter and Hansen, 2010, and a new species of *Stethacanthus* Newberry, 1889, *S. concavus* sp. nov. In addition, there occur a few long, monocuspid tooth-like denticles, similar to those observed in *Cobelodus* Zangerl, 1973, probably representing the head cover or the spine-brush complex. A review of the available information on the fossil record of Symmoriiformes has revealed that the group existed from the Late Devonian (Famennian) till the end of the Middle Permian (Capitanian).

Key words: Symmoriiformes; Microfossils; Carboniferous; Indian Cave Sandstone; USA Midcontinent.

INTRODUCTION

The Symmoriiformes (Symmoriida *sensu* Zangerl 1981) are a group of Palaeozoic cladodont sharks sharing several common characters: relatively short skulls, large eyes, terminal mouth, epicercal but externally symmetrical caudal fin, pectoral fins with a long metapterygial axis, posterior dorsal fin devoid of a fin-spine, anterior dorsal fin missing or transformed (at least in males) into a specific structure, such as a spine-brush complex in *Stethacanthus* Newberry, 1889 and *Orestiacanthus* Lund, 1984 or an anteriorly directed head spine in falcatids (see e.g., Zangerl 1981, 1984; Lund 1984, 1985, 1986; Williams 1985; Coates and Sequeira 2001; Maisey 2007, 2009). Their skin is generally naked, but specialised denticles usually occur on the head, on the spine, protecting the lateral line and there also occur very characteristic, complex, *Stemmatias*-type pharyngeal denticles.

The teeth of symmoriiforms are of cladodont design, i.e., with the central main cusp the highest, but they differ considerably within the group, as far as the

size and shape is concerned [compare the thick median cusp, almost a centimetre long, in *Stethacanthus neilsoni* (Traquair, 1898), and the minute, 0.5 mm wide, multicuspid, comb-like tooth of *Denaëa wangi* Wang, Jin and Wang, 2004; Ginter *et al.* 2010, figs 58A–C and 61, respectively]. However, again several important common dental features can be observed. First of all, the cusps are structurally separate (the feature first noted by Sequeira and Coates 2000, pp. 162, 167); only rarely, when the bases of cusps are very close to each other, a thin enameloid connection exists. Also, the cusps and base tissues are clearly separated and they react differently to diagenetic processes (see e.g., Ginter *et al.* 2010, fig. 58D). The cusps are rounded in cross section; the labio-lingual compression, if it occurs at all, is slight. The bases are usually flat, with a broad lingual extension, and in primitive members of the group (e.g., *Stethacanthus*) provided with pronounced articulation devices (single button + basolabial projection). In later taxa (e.g., *Stethacanthulus* Zangerl, 1990) such devices may be highly modified or are absent.

In 1974, Clair R. Ossian presented his Ph. D. dissertation, in which he described numerous fossils, mainly of vertebrate and botanic origin, from the locality at Peru, Nemaha County, Nebraska, USA (Ossian 1974). The fossils were found in the Indian Cave Sandstone, which is presumed to represent the Upper Pennsylvanian, Gzhelian (for the geological setting see Fischbein *et al.* 2009; Ginter 2016, pp. 300, 301, text-fig. 1). Among the very diverse vertebrate macro- and microfossils – counted in thousands – he reported about 20 chondrichthyan tooth-based taxa, together with chondrichthyan dermal denticles, spines, and putative calcified cartilage fragments. Although the fossils were found in sandstone matrix, it was possible for Ossian to recover them by chemical processing, using acetic acid. Many chondrichthyan teeth are beautifully preserved, displaying only a slight, superficial abrasion. Larger vertebrate skeletal fragments, however, are usually broken.

Among the Chondrichthyes, Ossian (1974) identified a few cladodontomorphs, at least two xenacanthiforms, euselachians (see Ginter 2016), various euchondrocephalans (orodonts?, petalodonts, eugeodontiforms, and holocephalans), and iniopterygians. Unfortunately, the dissertation has never been published and the specimens, except the xenacanthiforms (Johnson 1984, 1999), have never been formally described. Based on our current knowledge it can be stated that only one tooth illustrated by Ossian (1974, pl. 3, figs 7–9), as *Cladodus gracilis* Newberry and Worthen, 1966, belongs to the Symmoriiformes. However, he also illustrated a few symmoriiform denticles, as teeth or tooth-whorls of *Styptobasis aculeata* Cope, 1894 (= *Cobelodus aculeatus*). The identity of these elements will be discussed below. It should be noted that Ossian (1974) reported 221 teeth of “*C. gracilis*” and 275 elements of “*S. aculeata*”, but illustrated only single representatives of each form, so the diversity in his material can be much broader than that inferred from the photographs.

In the Carnegie Museum of Natural History in Pittsburgh, USA, there occurs a collection of chondrichthyan teeth, apparently from the same locality and horizon (SE of Peru, Nebraska, Indian Cave Sandstone) as that studied by Ossian. The teeth are perfectly prepared and, as in the case of Ossian’s material, most of them are only slightly abraded. The quality and diversity of these teeth is so high that some of them were illustrated in the Handbook of Paleichthyology, vol. 3D (Ginter *et al.* 2010, figs 3, 6) as examples of various tooth forms in Palaeozoic chondrichthyans. In the collection there occur teeth of two symmoriiform species (one of them new and rather unusual) and a few

denticles similar to those referred by Ossian (1974) to *S. aculeata*. The description of these microfossils and the short review of post-Mississippian symmoriiforms are the main purposes of this paper.

Institutional abbreviation: CM, Carnegie Museum of Natural History, Pittsburgh, USA.

SYSTEMATIC PALAEOLOGY

Class Chondrichthyes Huxley, 1880
 Subclass Elasmobranchii Bonaparte, 1838
 Order Symmoriiformes Zangerl, 1981
 Family Falcatidae Zangerl, 1990
 Genus *Danaea* Pruvost, 1922

TYPE SPECIES: *Danaea fournieri* Pruvost, 1922.

Danaea saltsmani Ginter and Hansen, 2010
 (Text-fig. 1)


1983. Subtype 021; Tway and Zidek, p. 416, fig. 38.

2010. *Danaea saltsmani* sp. nov.; Ginter and Hansen, pp. 32–34, fig. 2.

MATERIAL: Two teeth with the same collection number, CM 44548 (a and b); Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska, USA.

DESCRIPTION: The following description is based mostly on the two teeth illustrated herein, but also on the specimens from the other localities in the Pennsylvanian of the USA Midcontinent (Ginter and Hansen 2010). The tooth crown of *D. saltsmani* is characterised by the following combination of features: the crown comprises five to eight cusps; in seven-cuspid teeth, the second to the outermost pair of the lateral cusps is the highest; the central cusp is circular in cross section, and curves markedly lingual with only slight sigmoid flexure; the cristae on the cusps are delicate, subparallel, extend from the base to the tip and generally do not converge; the cristae are equally spaced both labially and lingually and the lateral carinae are absent.

The tooth base of *D. saltsmani* is triangular to trapezoidal. The base of the triangle forms a straight labial margin (this feature is particularly evident in smaller specimens of this species) and the distal part of the lingual extension forms the apex of the triangle (or a shorter edge of the trapezoid). The base does not



Text-fig. 1. *Denaea saltsmani* Ginter and Hansen, 2010, from the Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska. A-C – tooth CM 44548a, in labial, oral and oblique lateral views; D-G – tooth CM 44548b, in lateral, labial, lingual and oral views

have a distinct button, but only a vague thickening or hump on the oral-lingual side: the lingual edge of the central part of the base is developed into a raised, bulbous area, in the centre of which, on the lingual margin, is a prominent, circular lingual foramen. Another, slightly smaller foramen perforates the labial side of this thickening (Text-fig. 1B, C). Yet another large foramen is present in the centre, or slightly lingual of centre, on the aboral surface. On the labial margin of the aboral surface, beneath the central cusp, is an elongate (mesio-distally), slightly raised, weak, articular boss. The lateral edges of the tooth base are slightly downturned in some specimens.

It is notable that both teeth of *D. saltsmani* from Peru (Text-fig. 1) and the holotype from the Noble limestone in Ohio (Ginter and Hansen 2010, fig. 2A–D) are asymmetrical and have six cusps in the crown. Such a condition rarely occurs, if at all, in the other falcatids (compare e.g., *Denaea* cf. *fournieri* in Ginter *et al.* 2015, figs 10A–E, 12).

REMARKS: Of the Pennsylvanian Falcatidae, only the teeth of *D. saltsmani* appear to closely resemble those of the type species, *D.ournieri* from the

Viséan of Belgium. The combination of such features as the presence of (however weak) articular devices (oral-lingual thickening and basolabial projection) and the faint, even, subparallel cristation of the cusps, distinguish *D.ournieri* and *D. saltsmani* from *Stethacanthulus meccaensis* (Williams, 1985) and *S. decorus* (Ivanov, 1999). Isolated teeth of *Denaea* cf. *fournieri* from the upper Viséan of Poland differ from *D. saltsmani* in having the outermost pair of the lateral cusps being the largest in seven-cuspid specimens (Ginter *et al.* 2015). Thus far, this condition has not been definitely recognised in the type material of *D.ournieri* from Belgium.

All the illustrated teeth of *Ozarcus mapesae* Pradel, Maisey, Tafforeau, Mapes and Mallatt, 2014 are pentacuspoid and the orolingual region of their bases is hardly visible, so, although these teeth are definitely falcatid in character (Pradel *et al.* 2014), it is difficult to compare them to the forms mentioned in the previous paragraph. A similar problem concerns the falcatids from the upper Mississippian of Bear Gulch, Montana, USA (*Falcatus* Lund, 1985 and *Damocles* Lund, 1986); the complete morphology of their teeth is still unknown.

A very well preserved seven- or perhaps even eight-cuspid tooth of *D. saltsmani* (putatively one of 221 specimens) was described by Ossian (1974, pl. 3, figs 7–9). He referred to it as *Cladodus gracilis* Newberry and Worthen, 1866, and suggested that *C. springeri* St. John and Worthen, 1875 might have represented the lower jaw dentition of the same species. It seems, however, that both mentioned Mississippian species (*C. gracilis* and *C. springeri*) sufficiently differ from the Pennsylvanian *D. saltsmani* to keep the latter as a separate taxon.

OCCURRENCE: Carboniferous, Upper Pennsylvanian, Gzhelian, USA (SE Nebraska; this paper). *Denaëa saltsmani* appears to range through much of the Pennsylvanian marine sequence in the Appalachian Basin (Kentucky, Ohio and Pennsylvania, USA; Pottsville through Conemaugh Formations; Hansen 1986; Ginter and Hansen 2010) and similar teeth were reported by Tway and Zidek (1983) from 16 localities in Virgilian (Upper Pennsylvanian) rocks of Iowa, Kansas, Missouri, Nebraska and Oklahoma, USA.

Family Symmoriidae Dean, 1909
Genus *Stethacanthus* Newberry, 1889

TYPE SPECIES: *Physonemus altonensis* St. John and Worthen, 1875.

Stethacanthus concavus sp. nov.
(Text-fig. 2)

2010. “undescribed symmoriiform”; Ginter *et al.*, fig. 3F, G.

ETYMOLOGY: From the concave appearance of the basolabial region of the tooth.

HOLOTYPE: Tooth, CM 44550b, from the Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska, USA.

MATERIAL: Five teeth; Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska, USA.

DIAGNOSIS: Symmoriiform shark with the following combination of dental features: five-cuspid crown composed of a long and slender median cusp, lingually convex and slightly flattened labially, covered with subdued, dense, subparallel cristae up to three-fourth of its height, and two pairs of small lateral cusplets, not

higher than one-fourth of the median cusp; reniform to rectangular base with a short, crescentic button on the orolingual side, perforated in the middle by a foramen leading to the main basal canal, and a corresponding to it short, labially concave basolabial ridge.

DESCRIPTION: Among the small collection of teeth of *S. concavus* sp. nov. several secondary differences between the teeth can be observed. The mesio-distal dimension of the base varies from 3 to 5 mm. The median cusp can be longer and narrower, with almost parallel lateral sides (Text-fig. 2E), or shorter and triangular (Text-fig. 2B). The base outline can be reniform or rectangular with rounded angles. The basolabial articulation device forms a simple, curved ridge (Text-fig. 2A), or a complex horseshoe structure with two labial projections (Text-fig. 2D, E). However, in all the instances, the ornamentation of the cusps and their relative separation, as well as the concave basolabial region unify these teeth.


REMARKS: Of all the known symmoriiforms, the teeth of *S. concavus* sp. nov. are morphologically the closest to those of *Stethacanthus*: the size relation between the median and lateral cusps in the pentacuspid crown and the general outline of the base are the common features. However, the deep basolabial concavity and the curved button corresponding to it distinguish the new species from all hitherto described stethacanthids. Thus, it is possible that in future, when more specimens of this type are studied, the taxon will be translated to a new genus.

OCCURRENCE: Carboniferous, Upper Pennsylvanian, Gzhelian, USA (SE Nebraska).

Stethacanthus sp.
(Text-fig. 3)

MATERIAL: Two teeth; Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska, USA.


DESCRIPTION and REMARKS: The first tooth referred here to as *Stethacanthus* sp. is generally similar to *S. concavus* sp. nov. (Text fig. 3A–C). It does have a curved, weakly developed button, but the concavity in the basolabial region is much shallower than in the typical specimens. Moreover, the cristae appear to reach the apex of the median cusp, whereas in the typical teeth they disappear lower, leaving the tip of the cusp completely smooth.



Text-fig. 2. *Stethacanthus concavus* sp. nov. from the Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska. A-C – holotype, CM 44550b, in basal, labial and basal/labial views; D, E – tooth CM 44550c, in basal and labial views; F-H – large tooth, CM 44550a, in oblique labial, oral and lingual views

The second tooth, in addition to having the above mentioned characters, displays strong interconnection between the cusps. This latter feature is unusual

for the symmoriids, but perhaps reflects some special position in the dentition. Otherwise, the tooth is definitely stethacanthid.



Text-fig. 3. Teeth of *Stethacanthus* sp. from the Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska. A-C – CM 44550d, in oblique lingual, oral and labial views; D, E – CM 44550e, in labial and oral views

OCCURRENCE: Carboniferous, Upper Pennsylvanian, Gzhelian, USA (SE Nebraska).


SYMMORIIFORM HEAD DENTICLES

In the collection from Peru, Nebraska, housed at the Carnegie Museum, there occur three monocuspid denticles (Text-fig. 4) with interestingly shaped bases. In all cases the tips of the cusps are broken, but it can be approximated that the length of the largest denticle was greater than 3 mm. The cusps are curved (backwards?), elliptical in cross section, slightly laterally compressed. They are ornamented with sub-parallel cristae, sometimes bifurcating near the base. For the obvious reason, it is unknown whether the cristae reach the apex of a cusp.

The base in two of the denticles (Text-fig. 4A–E) has a three-fold appearance, with two small lateral lobes and one, larger, directed backwards. The base of the smallest denticle (Text-fig. 4F, G) has a simple, trapezoidal base with rounded angles. Numerous irregularly placed openings occur on the upper side of

the base, around the basal part of the cusp; the lower side of the base is devoid of any foramina.

The size, form and ornamentation of the cusps are extremely similar to those of the denticles of *Cobelodus aculeatus* (Zangerl and Case 1976, fig. 16, upper row), a symmoriid from Pennsylvanian black shales in central USA. However, the bases of the denticles in the drawings provided by the mentioned authors do not have any specific features, they are simply rounded portions of spongy tissue attached at the base of the cusps. They may have lost their form due to the unfavourable preservation (pyritisation). Zangerl and Case (1976), as well as Zangerl (1981, fig. 8E) suggested that such denticles are upper jaw dentition teeth because they were found near the palatoquadrate. M. Ginter and A. Ivanov independently suggested around 2003–2004 (unpublished communications) that the shape of these denticles, and especially the form of their bases, precludes their belonging to the dentition and that they are rather head denticles, similar to those called *Cladodus pattersoni* by Newberry (1875) and Williams (1985). This concept, that the denticles represent a cranial cap, was subsequently suggested by



Text-fig. 4. Symmoriiform head denticles? from the Upper Pennsylvanian, Gzhelian, Indian Cave Sandstone, SE of Peru, Nebraska. A, B – CM 44550f, in lateral and basal/posterior views. C–E. CM 44550g, in basal, posterior and lateral views; F, G – CM 44550h, in external and lateral views

Maisey (2007, p. 63), Ginter *et al.* (2010, pp. 60, 61) and Ivanov (2016, p. 720).

The typical, tricuspid dentition teeth of *Cobelodus* (Ginter *et al.* 2010, fig. 57; Ivanov 2016, figs 2H–J, 3A–C) have not been found in the material from Peru, despite the very large number of symmoriiform teeth occurring in Ossian's collection. Therefore it is likely that the monocuspid denticles are associated with the teeth of *S. concavus* sp. nov, the only larger symmoriid present at the Peru locality.

It is interesting that the morphology of these denticles resembles that of alleged monocuspid teeth of *Celtiberina maderi* Wang, 1993, an Early Devonian shark from Spain (Ginter *et al.* 2010, fig. 21). This is yet another argument supporting the idea expressed by Turner (2004, p. 88) and followed by Ginter *et al.* (2010, p. 28) that *Celtiberina* elements are actually modified dermal denticles, and not dentition teeth.

FOSSIL RECORD OF THE SYMMORIIFORMES

The Devonian record of symmoriiform sharks is sparse. Here and there falcetid-like delicate teeth have

been reported from the Famennian (e.g., Ginter 2000, fig. 7B; 2001, fig. 2A–E) and the only partly articulated specimens (*Stethacanthus*) come from the upper Famennian Cleveland Shale of Ohio (Williams 1985). The situation is much better in the Mississippian, and particularly in the Viséan-Serpukhovian. In addition to the numerous isolated teeth, characteristic denticles (e.g., *Stemmatias* branchial denticles) and spines from spine-brush complexes, found all over the world (e.g., Zidek 1993; Ginter *et al.* 2015), quite a few fairly complete skeletons have been described. The major source of these are the Serpukhovian Bear Gulch Beds in Montana (*Falcatus*, *Damocles*, *Orestiacanthus*; Lund 1984, 1985, 1986), the Upper Mississippian of Scotland (*Stethacanthus*-like forms “*Akmonistion*” and “*Guttarensis*”; Sequeira and Coates 2000; Coates and Sequeira 2001), and the Viséan Black Marble in Belgium (*Denaëa*; Fournier and Pruvost 1928; Maisey 2008). There is also a recently described, well preserved skull with associated teeth of a falcetid, *Ozarcus mapesae*, from the Upper Mississippian (Chesterian) of Arkansas (Pradel *et al.* 2014).

The known distribution of post-Mississippian symmoriiforms is restricted to the areas of USA,

Table 1. The Symmoriiformes from post-Mississippian rocks.

| Current name | Original name | Important references | Age, occurrence | Articulated |
|--|---|---|--|-------------|
| Symmoriidae | | | | |
| <i>Symmorium reniforme</i> Cope, 1893 | | Williams 1985; Ginter 2002 | Pennsylvanian, USA | yes |
| <i>Cobelodus aculeatus</i> (Cope, 1894) | <i>Styptobasis aculeata</i> | Zangerl and Case 1976; Ivanov 2013a, 2016 | Pennsylvanian, USA; Permian, Russia [?] | yes |
| <i>Stethacanthus altonensis</i> (St. John and Worthen, 1875) | <i>Physonemus altonensis</i> | Newberry 1889; Williams 1985; Zidek 1993 | Pennsylvanian, USA | yes |
| <i>Stethacanthus concavus</i> sp. nov. | | this paper | Pennsylvanian, USA | no |
| <i>Stethacanthus rosanovi</i> (Minikh, 1999) | <i>Ctenacanthus rosanovi</i> + <i>Pinegia grunti</i> | Minikh 2004, 2006; Ivanov and Lebedev 2014 | Permian, Russia | no |
| Falcatidae | | | | |
| <i>Denaea saltsmani</i> Ginter and Hansen, 2010 | | Tway and Zidek 1983; Hansen 1986 | Pennsylvanian, USA | no |
| <i>Stethacanthulus meccaensis</i> (Williams, 1985) | <i>Denaea meccaensis</i> + <i>Stethacanthulus longipeniculus</i> | Zangerl 1990, 1995; Ginter and Hansen 2010; Ivanov 2013b, Ivanov <i>et al.</i> 2013, 2015 | Lower Pennsylvanian, Uzbekistan; Pennsylvanian and Permian, USA | yes |
| <i>Stethacanthulus decorus</i> (Ivanov, 1999) | <i>Denaea decora</i> | Ivanov 2005, 2013b | Lower Pennsylvanian, Uzbekistan; Pennsylvanian, USA; Permian, Russia | no |
| Family indet. | | | | |
| “ <i>Cladodus</i> ” <i>acuminatus</i> Newberry, 1856 | | Newberry 1875 | Pennsylvanian, USA | no |
| “ <i>Cladodus</i> ” <i>fulleri</i> St. John and Worthen, 1875 | | | Pennsylvanian, USA | no |
| “ <i>Cladodus</i> ” <i>pandatus</i> St. John and Worthen, 1875 | | | Pennsylvanian, USA | no |
| <i>Kawichthys moodiei</i> Pradel, Taf-foreau, Maisey and Janvier, 2011 | | | Pennsylvanian, USA | yes |
| <i>Kunguroodus obliquus</i> [?] (Ivanov, 2005) | <i>Cobelodus obliquus</i> | Lebedev 2009; Ivanov 2016 | Permian, Russia and Kazakhstan | no |

Russia, and Central Asia. However, their diversity is still high in the Pennsylvanian (see Table 1). The flourishing Serpukhovian falcatids were replaced with *Denaea saltsmani* and at least two species of *Stethacanthulus*. The most durable symmoriid genus, *Stethacanthus*, is well documented from the Pennsylvanian black shales of Indiana and adjacent states. The same formation yielded two new larger sharks, *Symmorium* Cope, 1893 and *Cobelodus*. Two interesting skulls with denticulated dorsal crests (but no associated teeth) were reported from the Upper Pennsylvanian (Virgilian) of Kansas, under the name of *Kawichthys* (Pradel *et al.* 2011). The newly described teeth of *Stethacanthus concavus* sp. nov. from Nebraska and a few forms incorrectly referred to *Cladodus* by 19th century writers, and still requiring revision, add to the picture.

The record of symmoriiforms decreases considerably at the Carboniferous–Permian boundary. Thus far almost only isolated teeth and dermal denticles have been recovered from Permian rocks. Among these, only three undoubtedly symmoriiform species

have been identified: *Stethacanthulus meccaensis* (Williams, 1985) (Capitanian, Texas; Ivanov *et al.* 2013, fig. 4a, b; 2015, pl. 1, figs 1–9), *S. decorus* (Ivanov, 1999) (Asselian and Sakmarian, Urals; Ivanov 1999, pl. 7, fig. 12, pl. 8; 2005, fig. 3K–P) and *Pinegocaptus rosanovi* (Minikh, 1999) (Kazanian, mainly Roadian, northern European Russia; Minikh 2004, pl. 1, figs 1, 2; Ivanov and Lebedev 2014, pl. 9, figs 1–4). The latter species is represented by large, up to a few centimetres high *Stethacanthus*-like teeth with a very thick median cusp, covered with numerous dense cristae. Not all authors accept the validity of the generic name *Pinegocaptus* Minikh, 2006. For instance, Ivanov *et al.* (2009, p. 244) stated that it is a junior synonym of *Stethacanthus*. Even if so, the unusual size and ornamentation of the median cusp suggests that *S. rosanovi* is a separate species.

The only partly articulated fragment of a symmoriid skeleton found in the Permian thus far, is a skeleton of a pectoral fin recovered from Artinskian deposits of the Middle Urals. It was described by Ivanov (2013a) as belonging to *Cobelodus*.

Special attention should be given to the tooth-based genus *Kunguroodus* Ivanov, 2016. The beautifully preserved teeth from the Lower Permian of South Urals and Cisurals (Russia and Kazakhstan; Ivanov 2005, 2016; Lebedev 2009) were originally described as *Cobelodus obliquus* by Ivanov (2005), but later, after Ivanov's study of the original specimens of *Cobelodus*, the species was transferred to a new genus (Ivanov 2016) within the Symmoriiformes. However, its symmoriid affinity is at least doubtful. The teeth of *K. obliquus* resemble such falcatids as *Denaia williamsi* Ginter and Hansen, 2010 (e.g., Ginter *et al.* 2015, fig. 13E) to some extent, but its cusps are divergent, compressed labio-lingually and, moreover, connected with a carina. Owing to such characters I put *Kunguroodus* on the list of post-Mississippian symmoriiforms (Table 1), but only tentatively.

Judging from the available information, the Symmoriiformes did not persist into the Upper Permian. The latest occurrences (*S. meccaensis*) were reported from the Capitanian (uppermost Middle Permian) of the Apache Mountains (Ivanov *et al.* 2013) and Guadalupe Mountains (Ivanov *et al.* 2015), West Texas. However, it must be taken into account that a few teeth very similar to those of the falcatids, and designated as such by Guinot *et al.* (2013, fig. 2a–e) were found in the deep marine deposits of the Lower Cretaceous, Valanginian, of southern France.

CONCLUSIONS

In the last two decades information on the symmoriiform sharks has considerably grown. On the one hand, several old articulated specimens have been restudied (Coates and Sequeira 2001; Maisey 2007, 2008, 2009) and new ones, particularly neurocrania, were added to the data base (Pradel *et al.* 2011, 2014) thanks to the new techniques employed, such as computer tomography. On the other hand, intensive works on symmoriid microfossils, particularly teeth, has helped to solve taxonomic problems and misunderstandings from the past (e.g., Ginter 2002; Ginter *et al.* 2010; Ivanov 2016) and considerably broadened our knowledge on the distribution of symmoriiforms in time and space (numerous publications by A.O. Ivanov since 1999). Currently we know that this group has existed since the Famennian through to the Middle Permian. The studies on the Permian are so promising that perhaps they will show an increased upper range of these sharks, even if the report from the Valanginian (Guinot *et al.* 2013) is considered dubious.

The symmoriiform microfossils from the Indian Cave Sandstone body near Peru, Nebraska, described herein, are only a small contribution to that growing mass of knowledge. The sample is very small, only about ten elements altogether. Even so, this has permitted a new species to be erected (*Stethacanthus concavus* sp. nov.). However, the sample from the Carnegie Museum is only the tip of an iceberg. It is certain that a re-examination of the whole material collected by Ossian from the same locality and housed at the University of Texas at Austin would bring very interesting results, particularly as far as the intraspecific diversity is concerned.

Acknowledgments

I am grateful to Dr. Richard Lund (Carnegie Museum of Natural History, Pittsburgh) for driving me across Pennsylvania to Pittsburgh and for his assistance at the Carnegie Museum, and to Dr. Alexander O. Ivanov (Saint Petersburg University) for providing me with the literature on post-Mississippian symmoriiforms and for the insightful review of the manuscript. I also thank Dr. Gary D. Johnson (Southern Methodist University, Dallas) for his review, correction of the text and useful taxonomic suggestions. My research in USA in 2003 was financed by the Polish-American Fulbright Commission scholarship.

REFERENCES

- Bonaparte, C. 1832–41. Iconografia della Fauna Italica per le quattro Classi degli Animali Vertebrati. Tomo 3: Pesci, 266 p. Salviucci; Rome.
- Coates, M.L. and Sequeira, S.E.K. 2001. A new stethacanthid chondrichthyan from the Lower Carboniferous of Bearsden, Scotland. *Journal of Vertebrate Paleontology*, **21**, 438–459.
- Cope, E.D. 1893. On *Symmorium*, and the position of the Cladodont sharks. *American Naturalist*, **27**, 999–1001.
- Cope, E.D. 1894. New and little known Paleozoic and Mesozoic fishes. *Journal of the Academy of Natural Sciences of Philadelphia* (2nd series), **9**, 427–448.
- Dean, B. 1909. Studies on fossil fishes (sharks, chimaeroids and arthrodiras). *Memoir of the American Museum of Natural History*, **9**, 211–287.
- Fischbein, S.A., Fielding, C.R. and Joeckel, R.M. 2009. Fluvial-estuarine reinterpretation of large, isolated sandstone bodies in epicontinental cyclothems, Upper Pennsylvanian, northern Midcontinent, USA, and their significance for understanding late Paleozoic sea-level fluctuations. *Sedimentary Geology*, **216**, 15–28.
- Fournier, G. and Pruvost, P. 1928. Description des poissons elasmobranches du Marbre Noir de Denée. *Mémoires de la Société géologique du Nord*, **9**, 1–23.

- Ginter, M. 2000. Late Famennian pelagic shark assemblages. *Acta Geologica Polonica*, **50**, 369–386.
- Ginter, M. 2001. Chondrichthyan biofacies in the late Famennian of Utah and Nevada. *Journal of Vertebrate Paleontology*, **21**, 714–729.
- Ginter, M. 2002. Taxonomic notes on *Phoebodus heslerorum* and *Symmorium reniforme* (Chondrichthyes, Elasmobranchii). *Acta Palaeontologica Polonica*, **47**, 547–555.
- Ginter, M. 2016. The heterodonty in euselachian sharks from the Pennsylvanian of Nebraska. *Acta Geologica Polonica*, **66**, 200–312.
- Ginter, M., Duffin, C.J., Dean, M.T. and Korn, D. 2015. Late Viséan pelagic chondrichthyans from Northern Europe. *Acta Palaeontologica Polonica*, **60**, 899–922.
- Ginter, M., Hampe, O. and Duffin, C.J. 2010. Chondrichthyes. Paleozoic Elasmobranchii: Teeth. In: Schultze, H.-P. (Ed.), *Handbook of Paleichthyology 3D*, pp. 1–168. Friedrich Pfeil; München.
- Ginter, M. and Hansen, M. 2010. Teeth of the cladodont shark *Denaëa* from the Carboniferous of central North America. In: Nowakowski, D. (Ed.), *Morphology and systematics of fossil vertebrates*, pp. 29–44. DN Publisher; Wrocław.
- Guinot, G., Adnet, S., Cavin, L. and Cappetta, H. 2013. Cretaceous stem chondrichthyans survived the end-Permian mass extinction. *Nature Communications*, 1–8, doi: 10.1038/ncomms3669.
- Hansen, M.C. 1986. Microscopic chondrichthyan remains from Pennsylvanian marine rocks of Ohio and adjacent areas, 536 p. Unpublished PhD Thesis, Ohio State University; Columbus.
- Huxley, T.H. 1880. *A manual of the anatomy of vertebrated animals*, 431 p. D. Appleton; New York.
- Ivanov, A. 1999. Late Devonian – Early Permian chondrichthyans of the Russian Arctic. *Acta Geologica Polonica*, **49**, 267–285.
- Ivanov, A. 2005. Early Permian chondrichthyans of the Middle and South Urals. *Revista Brasileira de Paleontologia*, **8**, 127–138.
- Ivanov, A.O. 2013a. New findings of the endoskeleton of Early Permian sharks. In: *Palaeontological and geological monuments and collections: significance of museums for their study and preservation. Collection of scientific articles*, pp. 35–41. Geological Institute of RAS, Kungur Historical-Architecture and Art Museum; Kungur. [In Russian]
- Ivanov, A.O. 2013b. Chondrichthyans from the early/late Carboniferous boundary beds of the Gissar Mountains, Uzbekistan. In: Lucas, S.G., DiMichele, W.A., Barrick, J.E., Schneider, J.W. and Spielmann, J.A. (Eds), *The Carboniferous–Permian transition. New Mexico Museum of Natural History and Science, Bulletin*, **60**, 143–151.
- Ivanov, A.O. 2016. Chondrichthyans from the Lower Permian of Mechetlino, South Urals. *Bulletin of Geosciences*, **91**, 717–729.
- Ivanov, A.O. and Lebedev, O.A. 2014. Permian chondrichthyans of the Kanin Peninsula, Russia. *Paleontological Journal*, **48**, 1030–1043.
- Ivanov, A., Lucas, S.G. and Krainer, K. 2009. Pennsylvanian fishes from the Sandia Formation, Socorro County, New Mexico. In: Lueth, V., Lucas, S.G. and Chamberlin, R.M. (Eds), *Geology of the Chupadera Mesa. New Mexico Geological Society 60th Annual Fall Field Conference Guidebook*, pp. 243–248. New Mexico Geological Society; Socorro.
- Ivanov, A.O., Nestell, G.P. and Nestell, M.K. 2013. Fish assemblage from the Capitanian (Middle Permian) of the Apache Mountains, West Texas, USA. In: Lucas, S.G., DiMichele, W.A., Barrick, J.E., Schneider, J.W. and Spielmann, J.A. (Eds), *The Carboniferous–Permian transition. New Mexico Museum of Natural History and Science, Bulletin*, **60**, 152–160.
- Ivanov A.O., Nestell, M.K. and Nestell, G.P. 2015. Middle Permian fish microremains from the Early Capitanian of the Guadalupe Mountains, West Texas, USA. *Micropaleontology*, **61**, 301–312.
- Johnson, G.D. 1984. A new species of *Xenacanthoidii* (Chondrichthyes, Elasmobranchii) from the Late Pennsylvanian of Nebraska. In: Mengel, R.M. (Ed.), *Papers in Vertebrate Paleontology Honoring Robert Warren Wilson. Special Publication of Carnegie Museum of Natural History*, **9**, 178–186.
- Johnson, G.D. 1999. Dentitions of late Palaeozoic *Orthacanthus* species and new species of *?Xenacanthus* (Chondrichthyes: Xenacanthiformes) from North America. *Acta Geologica Polonica*, **49**, 215–266.
- Lebedev, O.A. 2009. A new specimen of *Helicoprion* Karpinsky, 1899 from Kazakhstani Cisurals and a new reconstruction of its tooth whorl position and function. *Acta Zoologica*, **90**, 171–182.
- Lund, R. 1984. On the spines of the Stethacanthidae (Chondrichthyes), with a description of a new genus from the Mississippian Bear Gulch Limestone. *Géobios*, **17**, 281–295.
- Lund, R. 1985. The morphology of *Falcatus falcatus* (St. John and Worthen), a Mississippian stethacanthid chondrichthyan from the Bear Gulch Limestone of Montana. *Journal of Vertebrate Paleontology*, **5**, 1–19.
- Lund, R. 1986. On *Damocles serratus*, nov. gen. et sp. (Elasmobranchii, Cladodontida) from the Upper Mississippian Bear Gulch Limestone of Montana. *Journal of Vertebrate Paleontology*, **6**, 12–19.
- Maisey, J.G. 2007. The braincase in Paleozoic symmoriiform and cladoselachian sharks. *Bulletin of the American Museum of Natural History*, **307**, 1–122.
- Maisey, J.G. 2008. Some observations on *Denaëa fourmieri* (Chondrichthyes, Symmoriiformes) from the Lower Carboniferous of Belgium. *Acta Geologica Polonica*, **58**, 185–190.
- Maisey, J.G. 2009. The spine-brush complex in symmoriiform sharks (Chondrichthyes; Symmoriiformes), with comments on dorsal fin modularity. *Journal of Vertebrate Paleontology*, **2**, 14–24.
- Minikh, A.V. 1999. New shark species of the *Ctenacanthus* Ag. genus from the Kazanian stage of the Upper Permian; the

- basin of the Pinega river. *Trudy Nauchno-issledovatel'skogo instituta geologii Saratovskogo Gosudarstvennogo Universiteta im. N.G. Chernyshevskogo, New series*, **1**, 133–136. [In Russian]
- Minikh, A.V. 2004. New elasmobranchians from the Ufimian and Kazanian stages of the Permian in the northern regions of European Russia. In: Ivanov, A.V. (Ed.), Questions of Upper Paleozoic and Mesozoic paleontology and stratigraphy (Memories of Galina Grigor'evna Poslavskaya). *Trudy Nauchno-issledovatel'skogo instituta geologii Saratovskogo Gosudarstvennogo Universiteta im. N.G. Chernyshevskogo, New series*, **16**, 128–132. [In Russian]
- Minikh, A.V. 2006. Phylum Chordata. In: Grunt, T.A. (Ed.), Upper Permian of Kanin Peninsula, 180–186. Nauka; Moscow. [In Russian]
- Newberry, J.S. 1856. Description of several new genera and species of fossil fishes, from the Carboniferous strata of Ohio. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **8**, 96–100.
- Newberry, J.S. 1875. Descriptions of fossil fishes. *Report of the Geological Survey of Ohio*, **2** (2), 1–64.
- Newberry, J.S. 1889. The Paleozoic fishes of North America. *U.S. Geological Survey, Monograph*, **16**, 1–340.
- Newberry, J.S. and Worthen, A.H. 1866. Descriptions of new species of Vertebrates, mainly from Subcarboniferous limestone and Coal Measures of Illinois. *Geological Survey of Illinois*, **2**, 9–134.
- Ossian, C.R. 1974. Paleontology, paleobotany and facies characteristics of a Pennsylvanian delta in Southeastern Nebraska, 20 + 393 p. Unpublished PhD Dissertation, University of Texas, Austin, available at: <https://www.academia.edu/1424478>.
- Pradel, A., Tafforeau, P., Maisey, J.G. and Janvier, P. 2011. A new Paleozoic Symmoriiformes (Chondrichthyes) from the Late Carboniferous of Kansas (USA) and cladistic analysis of early chondrichthyans. *PLoS ONE*, **6** (9), e24938. doi:10.1371/journal.pone.0024938.
- Pradel, A., Maisey, J.G., Tafforeau, P., Mapes, R.H. and Mallatt, J. 2014. A Palaeozoic shark with osteichthyan-like branchial arches. *Nature*, **509**, 608–611.
- Pruvost, P. 1922. Description de *Denaëa fournieri*, sélacien nouveau du Marbre noir de Denée. Part 2 of Fournier, G. and Pruvost, P. Découverte d'un poisson nouveau dans le Marbre noir de Denée. *Bulletin de Academie Royal de Bruxelles Serie 5*, **8**, 213–218.
- Sequeira, E.K. and Coates, M.I., 2000. Reassessment of 'Cladodus' *neilsoni* Traquair: a primitive shark from the Lower Carboniferous of East Kilbride, Scotland. *Palaeontology*, **43**, 153–172.
- St. John, O. and Worthen, A.H. 1875. Description of fossil fishes. *Geological Survey of Illinois, Paleontology*, **6**, 245–488.
- Traquair, R.H. 1898. On *Cladodus neilsoni* (Traquair), from the Carboniferous limestone of East Kilbride. *Transactions of the Geological Society of Glasgow*, **11**, 41–50.
- Turner, S. 2004. Early vertebrates: analysis from microfossil evidence. In: Arratia, G., Wilson, M.V.H. and Cloutier, R. (Eds.), Recent advances in the origin and early radiation of vertebrates. Honoring Hans-Peter Schultze, pp. 67–94. Friedrich Pfeil; München.
- Tway, L.E. and Zidek, J. 1983. Catalog of Late Pennsylvanian ichthyoliths, part II. *Journal of Vertebrate Paleontology*, **2**, 414–438.
- Wang, R. 1993. Taxonomie, Palökologie und Biostratigraphie der Mikroichthyolithen aus dem Unterdevon Keltiberiens, Spanien. *Courier Forschungsinstitut Senckenberg*, **161**, 1–205.
- Wang, N.-Z., Jin, F. and Wang, W. 2004. Early Carboniferous fishes (acanthodian, actinopterygians and Chondrichthyes) from the east sector of north Quilian Mountain, China – Carboniferous fish sequence from the east sector of north Quilian Mountain (1). *Vertebrata Palasiatica*, **42**, 89–110. [In Chinese with English summary]
- Williams, M.E. 1985. The Cladodont level sharks of the Pennsylvanian black shales of central North America. *Palaeontographica Abteilung A*, **190**, 83–158.
- Zangerl, R. 1973. Interrelationships of early chondrichthyans. In: Greenwood, P.H., Miles, R.S. and Patterson, C. (Eds.), Interrelationships of fishes. *Zoological Journal of the Linnean Society*, Supplement 1, **53**, 1–14.
- Zangerl, R. 1981. Chondrichthyes I: Paleozoic Elasmobranchii. In: Schultze, H.-P. (Ed.), Handbook of Paleichthyology, Volume 3A, pp. 1–115. Gustav Fischer; Stuttgart.
- Zangerl, R. 1984. On the microscopic anatomy and possible function of the spine-“brush” complex of *Stethacanthus* (Elasmobranchii: Symmoriida). *Journal of Vertebrate Paleontology*, **4**, 372–378.
- Zangerl, R. 1990. Two new stethacanthid sharks (Stethacanthidae, Symmoriida) from the Pennsylvanian of Indiana, U.S.A. *Palaeontographica Abteilung A*, **213**, 115–141.
- Zangerl, R. 1995. The problem of vast numbers of cladodont shark denticles in the Pennsylvanian Excello Shale of Pike County, Indiana. *Journal of Paleontology*, **69**, 556–563.
- Zangerl, R. and Case, G.R. 1976. *Cobelodus aculeatus* (Cope), an anacanthous shark from Pennsylvanian black shales of North America. *Palaeontographica Abteilung A*, **154**, 107–157.
- Zidek, J. 1993. A large stethacanthid shark (Elasmobranchii: Symmoriida) from the Mississippian of Oklahoma. *Oklahoma Geology Notes*, **53**, 4–15.

Manuscript submitted: 12th January 2018

Revised version accepted: 19th June 2018